

# Gamay de Bouze N

Wine grape variety.



## Origin

This variety is resulting of a mutation of Gamay.

## Use

Wine grape variety.

## Name of the variety in France

Gamay de Bouze

## Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

## Regulatory data

In France, Gamay de Bouze is officially listed in the "Catalogue of vine varieties" on the A list and classified. This variety is also listed in the catalogue of Bulgarie.

---

## Description elements

The identification is based on:

- the slightly bronze young leaves,
- the skin of the berries is practically green up until the veraison period and a colored pulp at maturity.

# Evolution of cultivated areas in France

Year	1958	1968	1998	2008	2018
ha	77	54	385	310	126

## Genetic profile

Microsatellite	VVS2	VMD5	VMD7	VMD27	VRZAG62	VRZAG79	VMD25	VMD28	VMD32
Allele 1	131	232	239	178	194	244	238	216	239
Allele 2	135	236	249	186	204	246	238	245	271

### Cultivation and agronomic skills

The aptitudes of this variety are close to those of other types of teinturier Gamay varieties or Gamay.

### Susceptibility to Diseases and Pests

Gamay de Bouze's behavior towards diseases and pests is the same as that of Gamay.

### Clonal selection in France

The two certified Gamay de Bouze clones carry the numbers 223 and 431. A conservatory with 30 or so clones was planted in 2006 in the wine-growing region of Rhône.

### Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

### Phenology

Bud burst: 1 day after Chasselas.

Grape maturity: early-season, 1 week after Chasselas.

### Technological potential

The bunches are small and the berries are small to medium in size. The wines produced are colored (less than Gamay Fréaux but much more than Gamay).

They are less fruity than Gamay wines.



Plantgrape, all rights reserved,  
 plantgrape.fr, UMT Géno-Vigne®  
 INRAE - IFV - L'Institut Agro Montpellier