

# Grenache N

Wine grape variety.



## Origin

This variety was originally from Spain and was introduced to France in the Middle Ages.

## Use

Wine grape variety.

## Name of the variety in France

Grenache

## Description elements

The identification is based on:

- the tip of the young shoot with a low density of prostrate hairs,
- the green and shiny young leaves,
- the shoots with green internodes,
- the adult leaves with teeth with straight sides, no anthocyanin coloration of veins and petioles, a very twisted smooth leaf blade, and on the lower side of the leaves, no or a very low density of erect and prostrate hairs,
- the yellow woody shoots,
- the round-shaped berries.

## Synonymy

In the European Union, Grenache is officially called by other names: Alicante (Italy), Cannonau (Italy), Tocal rosso (Italy), Garnacha Tinta (Spain), Grenache noir (Bulgaria, Cyprus, Croatia) and Grenache rouge (Greece). These synonyms are officially recognized in France regarding plant propagation material.

## Regulatory data

In France, Grenache is officially listed in the "Catalogue of vine varieties" on the A list and classified. This variety is also listed in the catalogues of other Member States of the European Union: Austria, Bulgaria, Croatia, Cyprus, Greece, Italy, Malta, Portugal and Spain.

# Evolution of cultivated areas in France

Year	1958	1968	1979	1988	1998	2008	2018
ha	24886	51931	77669	86715	91619	98644	82966

## Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	135	223	239	191	188	258	240	243	239
Allele 2	143	238	243	191	188	258	254	243	249

### Cultivation and agronomic skills

This variety is very vigorous (strong vine trunks), is sometimes sensitive to coulure and can have issues with lignification. Grenache has an erect bearing and must be pruned short in gobelet or cordon. It easily shows symptoms of magnesium deficiency. It is better adapted to slightly acidic, gravel or stony (pebble) terroirs rather than to very limestone white soil.


### Clonal selection in France

The twenty-four Grenache clones carry the numbers 70, 134, 135, 136, 137, 224, 287, 362, 432, 433, 434, 435, 513, 514, 515, 516, 517, 814, 1064, 1065, 1212, 1270, 1271 and 1272. A conservatory of more than 360 clones was planted in the Côtes du Rhône in 1998.

### Phenology

Bud burst: 5 days after Chasselas.  
Grape maturity: late-season, 4 weeks after Chasselas.

### Debourrement




### Technological potential

Grenache's bunches are medium to large and the berries are medium in size. It has a very high sugar accumulation potential but the color decreases quickly as the yields increase. Acidity is generally low. Grenache produces sweet wines and very structured, ample and aromatic great wines suited to ageing, provided that they are planted in quality terroirs and that the yields are perfectly managed.

### Susceptibility to Diseases and Pests

This variety is very sensitive to downy mildew and phomopsis and is rather sensitive to bacterial necrosis, grey rot and vine moths. On the other hand, it is not very susceptible to mites and resists well against, ungrafted, *Meloidogyne arenaria* nematodes in the sands along the coast.

## Maturité



## Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.



*Plantgrape, all rights reserved,  
plantgrape.fr, UMT Géno-Vigne®  
INRAE - IFV - L'Institut Agro Montpellier*