

Aramon gris G

Wine grape variety.



Origin

This variety corresponds to the grey mutation of Aramon.

Use

Wine grape variety.

Name of the variety in France

Aramon gris

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Aramon gris is officially listed in the "Catalogue of vine varieties" on the A list and classified.

Description elements

The description corresponds to that of Aramon, except for the berry skin color which is grey when ripe.

Evolution of cultivated areas in France

Year	1958	1968	1979	1988	1998	2008	2018
ha	1356	1145	387	125	111	41	15

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	131	232	239	178	188	240	240	227	249
Allele 2	141	232	243	191	196	260	254	259	249

Cultivation and agronomic skills

The aptitudes are comparable to those of Aramon.

Susceptibility to Diseases and Pests

Aramon gris has the same characteristics as Aramon.

Clonal selection in France

The only certified Aramon gris clone carries the number 1280.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

Phenology

Bud burst: 2 days after Chasselas.

Grape maturity: late-season, 4 weeks and a half after Chasselas.

Technological potential

The bunches are large and the berries are very large. Aramon gris only produces common white wines without any particular qualities.



*Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier*