

# Ignea Rs

Table grape variety.



## Origin

Ignea or 185 Pirovano was obtained by Angelo Pirovano (Italy) by crossbreeding 46 Pirovano and 2 Pirovano.

## Use

Table grape variety.

## Name of the variety in France

Ignea

## Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

## Regulatory data

In France, Ignea was officially listed in the "Catalogue of vine varieties" on the A list.

## Description elements

The identification is based on:

- the tip of the young shoot with a high density of prostrate hairs,
- the green young leaves with bronze spots,
- the circular entire adult leaves, with an open V-shaped petiole sinus, small teeth with straight sides, no anthocyanin coloration, a slightly revolute, finely blistered leaf blade, and on the lower side of the leaves, no or a very low density of erect hairs and a low density of prostrate hairs,
- broad ellipsoid berries.

# Evolution of cultivated areas in France

Year 2018

ha 0

## Genetic profile

| Microsatellite | VVS2 | VVMD5 | VVMD7 | VVMD27 | VRZAG62 | VRZAG79 | VVMD25 | VVMD28 | VVMD32 |
|----------------|------|-------|-------|--------|---------|---------|--------|--------|--------|
| Allele 1       | 131  | 229   | 247   | 182    | 192     | 240     | 248    | 233    | 239    |
| Allele 2       | 141  | 234   | 251   | 191    | 204     | 258     | 254    | 243    | 263    |

### Cultivation and agronomic skills

Ignea displays low to moderate vigor.

### Susceptibility to Diseases and Pests

Ignea is fairly susceptible to grey rot, powdery and downy mildew.

### Clonal selection in France

There is no certified clone for this variety yet.

### Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.

### Phenology

Bud burst: 10 days after Chasselas.

Grape maturity: mid-season, 3 weeks after Chasselas.

### Technological potential

The bunches are medium to large in size, not very compact. The berries also are medium to large with a thick skin, fleshy pulp and a simple flavor. Ignea has a good storage and transport capacity.



Plantgrape, all rights reserved,  
plantgrape.fr, UMT Géno-Vigne®  
INRAE - IFV - L'Institut Agro Montpellier