

Mondeuse blanche B

Wine grape variety.



Origin

This variety is originally from the Savoie region. It does not correspond to the white mutation of Mondeuse, it is a distinct variety.

Use

Wine grape variety.

Name of the variety in France

Mondeuse blanche

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Mondeuse blanche is officially listed in the "Catalogue of vine varieties" on the A list and classified.

Description elements

The identification is based on:

- the tip of the young shoot with a medium density of prostrate hairs,
- the bronzed young leaves,
- the circular, medium, light green adult leaves, with five lobes, V-shaped lateral sinuses, an open U-shaped petiole sinus, long teeth compared to their width at the base with convex sides, no anthocyanin coloration of veins, a slightly blistered leaf blade, slightly curly on the edges, and on the lower side of the leaves, no or a very low density of erect and prostrate hairs,
- the round-shaped berries.

Evolution of cultivated areas in France

Year	1958	1979	1988	1998	2008	2018
ha	1	21	7	5	6	8.1

Genetic profile

Microsatellite	VVS2	VMD5	VMD7	VMD27	VRZAG62	VRZAG79	VMD25	VMD28	VMD32
Allele 1	131	225	239	186	188	252	240	216	239
Allele 2	149	229	260	188	194	260	262	257	261

Cultivation and agronomic skills

Mondeuse blanche is a grape variety with a drooping bearing.

Susceptibility to Diseases and Pests

This variety is not very susceptible to grey rot.

Clonal selection in France

The only certified Mondeuse blanche clone carries the number 1214.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

Phenology

Bud burst: 5 days after Chasselas.
Grape maturity: late-season, 4 weeks after Chasselas.

Technological potential

The bunches and berries are medium in size. Mondeuse blanche produces ordinary and moderately acidic wines, with a fairly high alcohol degree (good sugar accumulation potential). Wines can be a little aromatic if the harvest is carried out at maturity.



*Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier*