


# Précoce Bousquet B

Wine grape variety.



## Origin

This variety was selected by M. Bousquet in the Tarn region and corresponds to the traditional Hungarian variety Honigler.

## Use

Wine grape variety.

## Name of the variety in France

Précoce Bousquet

## Synonymy

In the European Union, Précoce Bousquet is officially called by other names : Bela (Slovenia) and Honigler (Hungary). These synonyms are officially recognized in France regarding plant propagation material.

## Regulatory data

In France, Précoce Bousquet is officially listed in the "Catalogue of vine varieties" on the A list and classified. This variety is also listed on the catalogues of other Member States of the European Union : Hungary and Slovenia.

## Description elements

The identification is based on:

- the tip of the young shoot with a high density of prostrate hairs,
- the circular adult leaves, with three or five lobes, closed U-shaped lateral sinuses with slightly overlapping lobes, an open U-shaped petiole sinus, medium teeth compared to their width at the base with straight sides, no anthocyanin coloration of veins, an involute leaf blade, and on the lower side of the leaves, a medium to high density of erect hairs and a low density of prostrate hairs,
- round-shaped or slightly flat berries.

# Evolution of mother vine surfaces

| Year | 1958 | 1968 | 1979 | 1988 | 2000 | 2008 | 2018 |
|------|------|------|------|------|------|------|------|
| ha   | 33   | 35   | 23   | 24   | 12   | 8    | 2.8  |

## Genetic profile

| Microsatellite | VVS2 | VVMD5 | VVMD7 | VVMD27 | VRZAG62 | VRZAG79 | VVMD25 | VVMD28 | VVMD32 |
|----------------|------|-------|-------|--------|---------|---------|--------|--------|--------|
| Allele 1       | 141  | 223   | 243   | 178    | 188     | 252     | 240    | 247    | 247    |
| Allele 2       | 143  | 229   | 247   | 191    | 204     | 260     | 254    | 259    | 251    |

### Cultivation and agronomic skills

Précoce Bousquet has moderate fertility and should preferably be pruned long.

### Clonal selection in France

There is no certified clone for this variety yet.

### Phenology

Bud burst: 4 days after Chasselas.

Grape maturity: early-season, half a week after Chasselas.


### Technological potential

The bunches are medium to large and the berries are medium in size. Précoce Bousquet produces rather warm but neutral wines.


### Susceptibility to Diseases and Pests

Précoce Bousquet is susceptible to grey rot.

### Debourrement



### Maturité



### Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.

- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.



*Plantgrape, all rights reserved,  
plantgrape.fr, UMT Géno-Vigne®  
INRAE - IFV - L'Institut Agro Montpellier*