

Roublot B

Wine grape variety.



Origin

This variety is originally from the Yonne region and based on published genetic analyses, would be the result of the crossbreeding between Pinot and Gouais blanc.

Use

Wine grape variety.

Name of the variety in France

Roublot

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Roublot is officially listed in the "Catalogue of vine varieties" on the A list and classified.

Description elements

The identification is based on:

- the tip of the young shoot with a medium density of prostrate hairs,
- the yellow young leaves with bronze spots,
- the adult leaves with five or seven lobes (the central lobe is divided), a slightly open petiole sinus or with slightly overlapping lobes, medium to long teeth compared to their width at the base with straight sides, a moderate anthocyanin coloration of veins, a twisted, revolute and blistered leaf blade, and on the lower side of the leaves, a medium density of erect hairs,
- the broad ellipsoid berries.

Evolution of cultivated areas in France

Year 2018

ha 0

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	131	225	239	178	194	240	238	216	249
Allele 2	135	238	239	182	196	244	238	245	271

Cultivation and agronomic skills

Roublot has a semi-erect bearing. Its fertility is Good but irregular, which is why it must be pruned long.

Susceptibility to Diseases and Pests

Roublot is susceptible to powdery mildew and grey rot.

Clonal selection in France

There is no certified clone for this variety yet.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

Phenology

Bud burst: same as Chasselas.

Grape maturity: early-season, 1 week and a half after Chasselas.

Technological potential

The bunches and berries are small. Roublot produces fine white wines with a deep yellow color and certain aromatic typicity.



Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier