

Bachet N

Wine grape variety.



Origin

This variety is originally from the north east of France (Aube) and based on published genetic analyses it would result from the crossbreeding of Gouais blanc and Pinot.

Use

Wine grape variety.

Name of the variety in France

Bachet

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Bachet is officially listed in the "Catalogue of vine varieties" on the A list and classified.

Description elements

The identification is based on:

- the tip of the young shoot with a very high density of prostrate hairs,
- the shoots with red striped internodes,
- the entire adult leaves, with an open V or U-shaped petiole sinus, medium teeth with straight sides, a moderate anthocyanin coloration of veins, an involute, sometimes slightly revolute on the edges and finely blistered leaf blade, and on the lower side of the leaves, a low to medium density of prostrate hairs,
- ellipsoid berries.

Evolution of cultivated areas in France

Year	1958	1979	1988	2018
ha	1	4	1	0

Genetic profile

Microsatellite	VVS2	VMD5	VMD7	VMD27	VRZAG62	VRZAG79	VMD25	VMD28	VMD32
Allele 1	141	232	243	176	188	238	248	227	271
Allele 2	149	236	249	186	204	246	254	235	271

Cultivation and agronomic skills

This variety has good vigor and is fairly productive. It appears to resist well against winter frosts. On the other hand, it is sensitive to chlorosis, drought and root asphyxia.

Susceptibility to Diseases and Pests

Bachet is susceptible to the main diseases: downy mildew, powdery mildew and grey rot.

Clonal selection in France

There is no certified clone for this variety yet.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

Phenology

Bud burst: 3 days before Chasselas.
Grape maturity: early-season, 1 week to 1 week and a half after Chasselas.

Technological potential

The bunches and berries are small. Bachet produces fairly common wines, relatively alcoholic and acidic, with herbaceous flavor.



Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier