


Admirable de Courtiller B

Table grape variety.







Origin

This variety was obtained by Mr. Courtiller in the 19th century in the Saumur region. Based on genetic analyses carried out in Montpellier, it would result from the crossbreeding of Bicane and Chasselas.

Use

Table grape variety.

Name of the variety in France

Admirable de Courtiller

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Admirable de Courtiller is officially listed in the "Catalogue of vine varieties" on the A list.

Description elements

The identification is based on:

- the tip of the young shoot with a medium to high density of prostrate hairs,
- the green young leaves with bronze spots,
- the pentagonal adult leaves with five or seven lobes, a slightly open U-shaped petiole sinus or with slightly overlapping lobes, medium teeth with convex sides, no anthocyanin coloration of veins, a smooth, shiny leaf blade and on the lower side of the leaves, a medium density of erect hairs and, no or very low density of prostrate hairs,
- the berries that are ellipsoid or ovoid, and which can become slightly pink when overripe.

Evolution of cultivated areas in France

Year	1958	1968	1979	2000	2008	2018
ha	909	898	369	35	22	6.9

Genetic profile

MicrosatelliteVVS2		VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	131	223	239	182	188	252	240	233	239
Allele 2	135	234	243	191	194	258	254	267	239

Cultivation and agronomic skills

Admirable de Courtiller displays low to moderate vigor and has a semi-erect bearing. Its yield potential is moderate. It has a tendancy to produce a lot of grappillons and is sometimes sensitive to coulure and millerandage. It is preferable to grow Admirable de Courtiller on dry and well-exposed terrains.

Clonal selection in France

The only certified Admirable de Courtiller clone carries the number 1308.

Phenology

Bud burst: 2 days after Chasselas.

Grape maturity: early-season, 1 week after Chasselas.


Technological potential

Admirable de Courtiller's bunches are medium in size and not very compact. The berries are medium to large in size, simple-flavored with a thin to medium skin and a juicy pulp.


Susceptibility to Diseases and Pests

Admirable de Courtiller is not very sensitive to rot.

Debourrement



Maturité



Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.











Plantgrape, all rights reserved, plantgrape.fr, UMT Géno-Vigne® INRAE - IFV - L'Institut Agro Montpellier