


# Corbeau N

Wine grape variety.



## Origin

This variety is originally from the Savoie region.

## Use

Wine grape variety.

## Name of the variety in France

Corbeau

## Synonymy

In France, this variety can officially be called "Douce noire" regarding plant propagation material. In the European Union, Corbeau is officially called by other names: Serbina (Italia) and Sevilhão (Portugal).

## Regulatory data

In France, Corbeau is officially listed in the "Catalogue of vine varieties" on the A list and classified.

## Description elements

The identification is based on:

- the tip of the young shoot with a high density of prostrate hairs,
- the green young leaves with bronze spots,
- the shoots with a ribbed surface,
- the wedge-shaped or pentagonal adult leaves, entire or with three or five, shallow lateral sinuses, an open U-shaped petiole sinus, small to medium teeth with convex sides, a weak anthocyanin pigmentation of veins, a flat or slightly involute, finely blistered leaf blade, and on the lower side of the leaves, a low density of erect and prostrate hairs,
- the round-shaped or slightly ellipsoid berries.

# Evolution of cultivated areas in France

Year	1958	1968	1979	2018
ha	962	576	55	3.3

## Genetic profile

Microsatellite	VVS2	VVM5	VVM7	VVM27	VRZAG62	VRZAG79	VVM25	VVM28	VVM32
Allele 1	149	225	249	186	194	252	248	227	239
Allele 2	149	236	262	186	200	260	262	243	271

### Cultivation and agronomic skills

Corbeau is fairly vigorous, with a semi-erect bearing. It can be trellised and pruned long.

### Clonal selection in France

The only certified Corbeau clone carries the number 1367.

### Phenology

Bud burst: 1 day after Chasselas.  
Grape maturity: mid-season, 2 weeks and and a half after Chasselas.


### Technological potential

Corbeau's bunches are small to medium in size and the berries are medium in size. This variety produces not very acidic, not very tannic, well colored wines with a pleasant taste. The sugar accumulation potential of this variety is low to moderate.


### Susceptibility to Diseases and Pests

Corbeau is not very susceptible to powdery mildew and grey rot. It is however more sensitive to downy mildew and anthracnose.

### Debourrement



### Maturité



### Bibliographic references

- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - ©2023 Plantgrape, all rights reserved

Institut Agro Montpellier, Marseillan, France.

- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

---



---

*Plantgrape, all rights reserved,  
plantgrape.fr, UMT Géno-Vigne®  
INRAE - IFV - L'Institut Agro Montpellier*