

New

Artys B

Wine grape variety.



Origin

Artys was obtained by INRAE. This interspecific hybrid results from the crossbreeding of a descendant of *Muscadinia rotundifolia* and Bronner.

Use

Wine grape variety.

Name of the variety in France

Artys

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Artys is officially listed in the "Catalogue of vine varieties" since 2024, on the A list and classified.

Description elements

The identification is based on:

- the tip of the young shoot with a moderate density of prostrate hairs,
- the green young leaves with a moderate density of prostrate hairs,
- the green shoots,
- the medium-sized, pentagonal adult leaves, with five lobes, a petiole sinus with slightly overlapping lobes, medium teeth with straight or convex sides, no anthocyanin coloration of veins, a slightly blistered leaf blade, and on the lower side of the leaves, a low density of prostrate hairs, - the broad ellipsoid berries.

Genetic profile

	MicrosatelliteVVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	135	223	239	186	184	240	238	227	249
Allele 2	143	236	241	186	184	246	238	243	271

Cultivation and agronomic skills

Artys is moderately vigorous, moderately fertile, with a semi-erect bearing.

Clonal selection in France

The only certified Artys clone carries the number 1398.

Phenology

Bud burst: 3 days after Chasselas.
Grape maturity: early-season, 2 weeks after Chasselas.

Debourrement

Technological potential


Artys' bunches are medium in size and moderately or not very compact. The berries are also medium in size and simple-flavored.

This variety produces neutral wines with fairly strong color intensity and a high acidity level. It can be used to make sparkling wines.


Susceptibility to Diseases and Pests

Artys is resistant to downy mildew and powdery mildew.

According to the breeder, based on current knowledge, two fungicide treatments for downy mildew and powdery mildew are essential to preserve resistance factors. This protection should be increased in case of high disease pressure.



Maturité



Bibliographic references

- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- [Artys variety note](#). INRAE Grand-Est Colmar, 2026.



*Plantgrape, all rights reserved,
plantgrape.fr, UMT Géo-Vigne®
INRAE - IFV - L'Institut Agro Montpellier*