


Danam B

Table grape variety.



Origin

This variety was obtained in 1958 by INRA by crossbreeding Dabouki and Muscat de Hambourg.

Use

Table grape variety.

Name of the variety in France

Danam

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Danam is officially listed in the "Catalogue of vine varieties" on the A list. This variety is also listed in the catalogue of Portugal.

Description elements

The identification is based on:

- the tip of the young shoot with a low to medium density of prostrate hairs,
- the green young leaves with bronze spots,
- the circular adult leaves, with five lobes, an open or with parallel edges U-shaped petiole sinus, long teeth with straight sides, no or a weak anthocyanin coloration of veins, a strongly blistered twisted leaf blade, and on the lower side of the leaves, no or a very low density of erect and prostrate hairs,
- the broad ellipsoid berries.

Evolution of cultivated areas in France

Year	1988	2000	2018
ha	3	1	0

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	147	229	249	180	186	240	244	235	271
Allele 2	149	232	249	182	188	248	254	257	271

Cultivation and agronomic skills

Danam displays moderate to strong vigor and is generally pruned short. It is not very affected by coulure and millerandage.

Susceptibility to Diseases and Pests

Danam is not very susceptible to rot.

Clonal selection in France

The only certified Danam clone carries the number 498.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.

Phenology

Bud burst: 5 days after Chasselas.
Grape maturity: mid-season, 3 weeks after Chasselas.

Technological potential

The bunches are very large. The berries are also very large with a thick skin and a simple-flavored pulp. The yield potential is high and it has a good transport capacity.



*Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier*